



FORM PTO-1390 (Rev. 10-95)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER MERCK 2032A
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. §371			U.S. APPLICATION NO. (If known, see 37 CFR §1.5) 10/089167
INTERNATIONAL APPLICATION NO. PCT/EP00/08939	INTERNATIONAL FILING DATE 13 SEPTEMBER 2000	PRIORITY DATE CLAIMED 28 SEPTEMBER 1999	
TITLE OF INVENTION QUINAZOLINONES			
APPLICANT(S) FOR DO/EO/US MEDERSKI, Werner, et al.			
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:</p> <ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. §371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. §371. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. §371(c)(2)) <ol style="list-style-type: none"> <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input checked="" type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. §371(c)(2)). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. §371(c)(3)). <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)). <p>Items 11. to 16. below concern document(s) or information included:</p> <ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 C.F.R. §§1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. §§3.28 and 3.31 is included. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input type="checkbox"/> A change of power of attorney and/or address letter. <input type="checkbox"/> Other items or information: 			

U.S. APPLICATION NO. (if known, see 37 CFR §1.5) 10/089167		INTERNATIONAL APPLICATION NO. PCT/EP00/08939		ATTORNEY'S DOCKET NUMBER MERCK 2032A	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR §1.492 (a) (1) - (5)): Search Report has been prepared by the EPO or JPO..... \$890.00 International preliminary examination fee paid to USPTO (37 CFR §1.482)..... \$710.00 No international preliminary examination fee paid to USPTO (37 CFR §1.482) but international search fee paid to USPTO (37 CFR §1.445(a)(2))..... \$740.00 Neither international preliminary examination fee (37 CFR §1.482) nor international search fee (37 CFR §1.445(a)(2)) paid to USPTO..... \$1040.00 International preliminary examination fee paid to USPTO (37 CFR §1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT = \$890.00				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 C.F.R. §1.492(e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	9 - 20 =	0	x \$ 18.00	\$0.00	
Independent claims	1 - 3 =	0	x \$ 84.00	\$0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$ 280.00		
TOTAL OF ABOVE CALCULATIONS =				\$890.00	
Reduction of 1/2 for filing by small entity, if applicable. A Verified Small Entity Statement must also be					
SUBTOTAL =				\$890.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 C.F.R. §1.492(f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
TOTAL NATIONAL FEE =				\$890.00	
Fee for recording the enclosed assignment (37 C.F.R. §1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. §§3.28, 3.31). \$40.00 per property.					
TOTAL FEES ENCLOSED =				\$890.00	
				Amount to be refunded:	
				charged:	
a. <input checked="" type="checkbox"/> A check in the amount of <u>\$890.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. <u>13-3402</u> in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-3402</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 C.F.R. §§1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. §1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Customer Number 23,599					
 23599 PATENT TRADEMARK OFFICE			SIGNATURE  Anthony J. Zelano NAME <u>27,969</u> REGISTRATION NUMBER		
Filed: 27 MARCH 2002 AJZ:kmo					



5000

10089167.082902

JC02 Rec'd PCT/PTO 22 MAY 2002

PC7

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of :

W. MEDERSKI et al.

Group Art Unit: TBA

Serial No.: 10/089,167

Examiner: TBA

Filed: March 27, 2002

For: QUINAZOLINONES

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Prior to initial examination, please amend the above-identified application as follows:

IN THE SPECIFICATION:

Please insert on page 1, after the title:

--This application claims priority of provisional application U.S. Serial No. 60/287,586,
filed September 28, 1999. --


MERCK 2032 A

REMARKS

The amendment to the specification clarifies that the present application claims priority to 60/325,777.

The Commissioner is hereby authorized to charge any fees associated with this response or credit any overpayment to Deposit Account No. 13-3402.

Respectfully submitted,



Nancy J. Axelrod (Reg. 44,014)
Attorney/Agent for Applicant(s)


MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
Arlington Courthouse Plaza 1, Suite 1400
2200 Clarendon Boulevard
Arlington, Virginia 22201
Telephone: (703) 243-6333
Facsimile: (703) 243-6410

Filed: May 22, 2002

K:\Merck\2033\A\PRE AMD.doc

Quinazolinones

The invention relates to substituted quinazolinones of the formula I



in which

- 10 R and R' are independently of each other H, A, OH, OA, OCH₂-Ar, Hal, NH₂, NHA, NA₂, NO₂, CN, C(O)R², CONH₂, CONHA, CONA₂, COOH, COOA or SO₂A,
- R² and R³ are independently of each other H, A, -C(=NH)-NH₂ or solid phase,
- 15 R⁴ is Ar, phenylalkyl, cycloalkyl or Het,
- Y may be absent and, if present, is alkenyl having 2 to 4 carbon atoms,
- A is unbranched or branched alkyl having 1 to 6 carbon atoms,
- Ar is phenyl, naphthyl, biphenyl or benzofuranyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA,
- 20 CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂,
- Het is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and
- 25 the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, NH₂, NHA, NA₂, COOH, COOA, phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂ or thiophenyl which is
- 30 unsubstituted or mono-, di- or trisubstituted by A, OH, OA,

CF_3 , OCF_3 , Hal, CN, COOH, COOA, NH_2 , NHA, NA_2 , NO_2 ,
 SO_2NH_2 , SO_2NAH or SO_2NA_2 ,

Hal is F, Cl, Br or I,
n is 0, 1, 2 or 3,
5 m is 0, 1, 2 or 3,
and their pharmaceutically tolerable salts and solvates.

Similar compounds having a quinazolinone parent structure as a
combinatorial library are disclosed in WO 98/11438. W.D. Dean et al, J.
10 Het. Chem. 1982, 1117-24 and L. Legrand et al, Bull. Soc. Chim. Fr. 1976,
1853-6 describes methods for the synthesis of similar quinazolinone
compounds.

The invention is based on the object of finding novel compounds having
15 valuable properties, in particular those which can be used for the
production of medicaments.

It has been found that the compounds of the formula I and their salts or
solvates have very valuable pharmacological properties together with good
tolerability.

20 They act especially as GPIbIX inhibitors, in particular inhibiting the
interaction of this receptor with the ligand von Willebrand factor (vWF). This
action can be demonstrated, for example, by a method which is described
by S. Meyer et al. in J. Biol. Chem. 1993, 268, 20555-20562. The property
as GPIbIX alpha-thrombin receptor (N.J. Greco, Biochemistry 1996, 35,
25 915-921) can also be blocked by the compounds mentioned.

The significance of GPIbIX as an adhesion receptor on platelets, which
mediates the primary interaction of platelets with an arteriosclerotically
modified vascular wall via binding to the vWF expressed there, has been
30 described by many authors (e.g. Z.M. Ruggeri in Thromb. Hemost. 1997,

78, 611-616). The activation of another platelet adhesion receptor, GPIIb/IIIa, following the GPIbIX-vWF interaction, leads to platelet aggregation and thus to thrombotic vascular occlusion.

- 5 A GPIbIX antagonist can thus prevent the start of thrombus formation and thus also release of active substances from the platelets which, for example, promote thrombus growth and have an additional trophic action on the vascular wall. This has been shown with inhibitory peptides or antibodies in various experimental models (e.g. H Yamamoto et al.,
10 Thromb. Hemost. 1998, 79, 202-210).

In the case of higher shear forces, the blocking action of GPIbIX inhibitors exerts its maximum effect, as described by J.J. Sixma et al. in Arteriosclerosis, Thrombosis, and Vascular Biology 1996, 16, 64-71.

- 15 According to the flow chamber method used there, the compounds of the formula I can be characterized as GPIbIX inhibitors in whole blood.

- The inhibition of thrombus formation of the GPIbIX inhibitors can be measured by a modified Born method (Nature 1962, 4832, 927-929) using
20 botrocetin or ristocetin as an aggregation stimulant.

- The compounds of the formula I according to the invention can therefore be employed as pharmaceutical active compounds in human and veterinary medicine. They act as adhesion receptor antagonists, in particular as
25 glycoprotein IbIX antagonists, and are suitable for the prophylaxis and/or therapy of thrombotic disorders and sequelae deriving therefrom. The preferentially best action is to be expected in the case of thrombotic disorders in the arterial vascular system, but GPIbIX inhibitors also have an effect in the case of thrombotic disorders in the venous vascular bed. The
30 disorders are acute coronary syndromes, angina pectoris, myocardial

infarct, peripheral circulatory disorders, stroke, transient ischaemic attacks, arteriosclerosis, reocclusion/restenosis after angioplasty/stent implantation. The compounds can furthermore be employed as anti-adhesive substances where the body comes into contact with foreign surfaces such as implants, catheters or cardiac pacemakers.


Therefore, the invention relates further to compounds of the formula I according to Claim 1 and their physiologically acceptable salts or solvates as pharmaceutical active compounds.

The invention relates to compounds of the formula I according to Claim 1 and their physiologically acceptable salts or solvates as glycoprotein IblX antagonists.

Comparison medication introduced onto the market which may be mentioned are aspirin and GPIIb/IIIa antagonists.


The invention relates to the compounds of the formula I and their salts or solvates, and to a process for the preparation of these compounds and their salts or solvates, characterized in that

- a) a compound of the formula I is liberated from one of its functional derivatives by treating with a solvolysing or hydrogenolysing agent, or
- b) in stage 1) a compound of the formula II




in which
X is Cl, Br, OH or a reactive esterified OH group and

Q is NH_2 or NHA , either of which is optionally protected, and R and R^1 are optionally protected when they are or contain NH_2 or NHA , is reacted with a compound of the formula III




in which R^2 , R^3 , n and m have the meanings indicated in Claim 1, to give a compound of formula IV



in which R , R^1 , R^2 , R^3 , Q , n and m have the meanings indicated above, and

in stage 2) a compound of formula IV as indicated above is if necessary deprotected to give a compound of formula IV in which Q is NH_2 or NHA and is reacted with a compound of formula V



in which R^4 and Y have the meanings indicated in Claim 1,

or

c) a radical R , R^1 , R^2 , R^3 and/or R^4 is converted into another radical R , R^1 , R^2 , R^3 and/or R^4 by, for example

- converting an amino group into a guanidino group by reaction with an amidinating agent,
- reducing a nitro group, sulfonyl group or sulfoxyl group,
- etherifying an OH group or subjecting an OA group to ether cleavage,
- alkylating a primary or secondary amino group,
- partially or completely hydrolysing a CN group,
- cleaving an ester group or esterifying a carboxylic acid radical,

- reacting an aryl bromide, aryl iodide, heteroaryl bromide or heteroaryliodide to give the corresponding coupling products by means of a Suzuki coupling with boronic acids,
 - or carrying out a nucleophilic or electrophilic substitution,
- 5 and/or
- a base or acid of the formula I is converted into one of its salts or solvates.

The compounds of the formula I can have a chiral centre and therefore occur in a number of stereoisomeric forms. All these forms (e.g. R and S

10 forms) and their mixtures (e.g. the RS forms) are included in the formula I.

The compounds according to the invention also include so-called prodrug derivatives, i.e. compounds of the formula I modified with, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved

15 in the body to give the active compounds according to the invention.

Furthermore, free amino groups as substituents of compounds of the formula I can be provided with appropriate conventional protective groups. Solvates of the compounds of the formula I are understood as meaning

20 adducts of inert solvent molecules to the compounds of the formula I which are formed on account of their mutual power of attraction. Solvates are, for example, mono- or dihydrates or alcoholates.

The abbreviations used have the following meanings:

- 25 BOC tert-butoxycarbonyl,
CBZ benzyloxycarbonyl,
DBU 1,8-diazabicyclo[5.4.0]undec-7-ene,
DCC dicyclohexylcarbodiimide,
DCE dichloroethane,
30 DDQ 2,3-dichloro-5,6-dicyano-1,4-benzoquinone,

- DMA dimethylacetamide,
 DMF dimethylformamide,
 dppf 1,1'-bis(diphenylphosphino)ferrocene,
 Et ethyl,
 5 Fmoc fluorenylmethoxycarbonyl,
 HBTU O-(benzotriazolyl)-N,N,N',N'-tetramethyluronium hexafluoro
 phosphate,
 Me methyl,
 Mtr 4-methoxy-2,3,6-trimethylphenylsulfonyl,
 10 OBut tert-butyl ester,
 OMe methyl ester,
 OEt ethyl ester,
 POA phenoxyacetyl,
 Ph phenyl,
 15 TEA triethylamine,
 TFA trifluoroacetic acid.

- In the above formulae, A is alkyl and has 1 to 6, preferably 1, 2, 3 or 4 C
 atoms. Alkyl is preferably methyl, furthermore ethyl, propyl, isopropyl, butyl,
 20 isobutyl, sec-butyl or tert-butyl, additionally also pentyl, 1-, 2- or
 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-,
 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or
 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or
 1,2,2-trimethylpropyl.
 25 A is preferentially methyl.

Alkenyl having 2 to 4 carbon atoms is preferably vinyl or buta-1,3-dienyl;
 vinyl is particularly preferred.

Ar is phenyl, naphthyl, biphenyl or benzofuranyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂.

- 5 Ar is preferentially phenyl, preferably - as indicated - mono- di- or trisubstituted phenyl, specifically preferentially phenyl, 2-, 3- or 4-methylphenyl, 2-, 3- or 4-ethylphenyl, 2-, 3- or 4-propylphenyl, 2-, 3- or 4-isopropylphenyl, 2-, 3- or 4-tert-butylphenyl, 2-, 3- or 4-aminophenyl, 2-, 3- or 4-N,N-dimethylaminophenyl, 2-, 3- or 4-sulfonamidophenyl, 2-, 3- or 4-nitrophenyl, 2-, 3- or 4-hydroxyphenyl, 2-, 3- or 4-methoxyphenyl, 2-, 3- or 4-ethoxyphenyl, 2-, 3- or 4-trifluoromethylphenyl, 2-, 3- or 4-trifluoromethoxyphenyl, 2-, 3- or 4-carboxyphenyl, 2-, 3- or 4-cyanophenyl, 2-, 3- or 4-fluorophenyl, 2-, 3- or 4-chlorophenyl, 2-, 3- or 4-bromophenyl. Furthermore Ar is preferentially unsubstituted naphthyl, biphenyl or
- 15 benzofuran-5-yl.
- Phenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 4-tert-butylphenyl, 4-dimethylaminophenyl, 4-methoxyphenyl, 3-methoxyphenyl, 3-chlorophenyl, 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 3',5'-dimethoxybiphenyl-4-yl, 2',4'-dimethoxybiphenyl-4-yl, biphenyl-4-yl, naphthalen-1-yl, naphthalen-2-yl or benzofuran-5-yl is
- 20 particularly preferred for Ar.

- Cycloalkyl preferably has 3-7 C atoms and is preferably cyclopropyl or cyclobutyl, furthermore preferably cyclopentyl or cyclohexyl, and further
- 25 also cycloheptyl; cyclohexyl is particularly preferred .

Hal is preferably F, Cl or Br.

- Het is a saturated, partially or completely unsaturated mono- or bicyclic
- 30 heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1

or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, NH₂, NHA, NA₂, COOH, COOA, phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂,
 5 SO₂NH₂, SO₂NAH or SO₂NA₂ or thiophenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂.

Het is preferably substituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂ or thiophenyl
 10 which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂ or unsubstituted 2- or 3-furyl, 2- or 3-thiophenyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or
 15 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -4- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl,
 20 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-1H-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or
 25 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl,

- 3-, 4-, 5-, 6-, 7- or 8-cinnoliny, 2-, 4-, 5-, 6-, 7- or 8-quinazoliny. The heterocyclic radicals can also be partially or completely hydrogenated. Het can thus also be 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or -5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl, 5 tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -3-pyrrolyl, tetrahydro-1-, -2- or 4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4-, -5-, -6-, -7-1H-indolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3- 10 or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1,2,3,6-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 1-, 2-, 3- or 4-azepanyl, 2-, 3- or 4-morpholiny, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or 15 -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinoliny, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinoliny.
- 20 2-Furyl, thiophen-2-yl, thiophen-3-yl, 5-(3,4-dimethoxyphenyl)-thiophen-2-yl or 5-[2,2']bithiophenyl is particularly preferred for Het.

Phenylalkyl preferably has 7, 8, 9 or 10 carbon atoms and is preferably phenylmethyl, phenylethyl, phenylpropyl or phenylbutyl; phenylethyl is 25 particularly preferred.

The term solid phase indicates a resin for solid-phase chemistry, especially for combinatorial chemistry, i.e. by robot- and computer-assisted syntheses, and subjected to mass screening as indicated in US 5,463,564; 30 M. A. Gallop et al., J. Med. Chem. 1994, 37, 1233-1251 and 1385-1401

and M.J. Sofia, Drugs Discovery Today 1996, 1, 27-34). The polymeric material of the solid phase is generally chosen from the group consisting of cross-linked polystyrene, cross-linked polyacrylamide or other resins, natural polymers or silicagels.

5

The group of cross-linked polystyrene, cross-linked polyacrylamide or other resins includes e.g. polyacrylamide, polymethacrylamide, polyhydroxyethylmethacrylate, polyamide, polystyrene, (meth)acrylate copolymers, for instance from (meth)acrylic acid, esters of (meth)acrylic acid and/or 2-methylene-succinic acid, but-2-enoic acid or maleic acid, polyurethanes or other copolymers.

10

Suitable terminal functional groups or linkers on the surface of the resin have to be chosen to attach the compounds to the resin. There exists a variety of commercially available resins, e.g. in Novabiochem - The Combinatorial Chemistry Catalog, March 99. Examples for suitable resins are carbonate resins with a modified carbonate group as terminal functional group like p-nitrophenylcarbonate resin, halogenated resins like Merrifield resin (chloromethylpolystyrene) or carboxy resins like carboxy polystyrene resin or NovaSyn® TG Carboxy Resin. p-Nitrophenylcarbonate resin is particularly preferred. These and other types of resins well known in the art can be used in the subject invention.

15

20

R and R¹ are independently of each other H, A, OH, OA, OCH₂-Ar, Hal, NH₂, NHA, NA₂, NO₂, CN, C(O)R², CONH₂, CONHA, CONA₂, COOH, COOA or SO₂A, where A, Ar, Hal have a preferred meaning indicated beforehand and R² has a preferred meaning indicated in the following.

25

R is preferentially H.

30

R¹ is preferentially H, A, OA or Hal.

The preferred position of R^1 is the 6- or 7-position of the quinazolinone ring system.

5 R^2 and R^3 are independently of each other H, A, $-C(=NH)-NH_2$ or a solid phase, where A or solid phase have a preferred meaning indicated beforehand.

R^2 is preferentially H.

10 R^3 is preferentially H or $-C(=NH)-NH_2$, particularly preferred is H.

R^4 is Ar, phenylalkyl, cycloalkyl or Het, where Ar, phenylalkyl, cycloalkyl or Het have a preferred meaning indicated beforehand. R^4 is preferentially phenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 4-tert-butylphenyl, 4-dimethylaminophenyl, 4-methoxyphenyl, 3-methoxyphenyl, 15 3-chlorophenyl, 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 3',5'-dimethoxybiphenyl-4-yl, 2',4'-dimethoxybiphenyl-4-yl, biphenyl-4-yl, naphthalen-1-yl, naphthalen-2-yl or benzofuran-5-yl, phenylethyl, cyclohexyl, 2-furyl, thiophen-2-yl, thiophen-3-yl, 5-(3,4-dimethoxyphenyl)-thiophen-2-yl or 5-[2,2']bithiophenyl.

20 Y may be absent and, if present, is alkenyl having 2 to 4 carbon atoms. Y is preferentially absent or vinyl.

25 n and m are each independently of each other 0, 1, 2 or 3, particularly preferred 1.

Some preferred groups of compounds can be expressed by the following subformulae Ia to Im, which correspond to the formula I and in which the radicals not designated in greater detail have the meanings indicated in 30 formula I, but in which

- in Ia R is H and
R¹ is H, A, OA or Hal;
- 5 in Ib R is H,
R¹ is H, A, OA or Hal and
Y is absent;
- 10 in Ic R is H,
R¹ is H, A, OA or Hal and
Y is alkenyl having 2 to 4 carbon atoms;
- 15 in Id R is H,
R¹ is H, A, OA or Hal,
R² is H and
R⁴ is Ar;
- 20 in Ie R is H,
R¹ is H, A, OA or Hal,
R² is H and
R⁴ is phenylalkyl;
- 25 in If R is H,
R¹ is H, A, OA or Hal,
R² is H and
R⁴ is cycloalkyl;
- 30 in Ig R is H,
R¹ is H, A, OA or Hal,
R² is H and

- R⁴ is Het;
- in lh R is H,
 R¹ is H, A, OA or Hal,
 5 R² is H,
 R³ is H,
 R⁴ is phenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 4-tert-butylphenyl, 4-dimethylaminophenyl, 4-methoxyphenyl, 3-methoxyphenyl, 3-chlorophenyl, 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 3',5'-dimethoxybiphenyl-4-yl, 2',4'-dimethoxybiphenyl-4-yl, biphenyl-4-yl, naphthalen-1-yl, naphthalen-2-yl or benzofuran-5-yl,
 10 phenylethyl, cyclohexyl, 2-furyl, thiophen-2-yl, thiophen-3-yl, 5-(3,4-dimethoxyphenyl)-thiophen-2-yl or 5-[2,2']bithiophenyl,
- 15 n is 1 and
 m is 1;
- in lk R is H,
 R¹ is H, A, OA or Hal,
 20 R² is H,
 R³ is H,
 Y is -CH=CH-,
 R⁴ is phenyl, 4-dimethylaminophenyl or 2,5-dimethoxyphenyl,
 n is 1 and
 25 m is 1;
- in lm R is H,
 R¹ is H, A, OA or Hal,
 R² is H,
 30 R³ is H,

- Y is absent,
- R⁴ is phenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 4-tert-butylphenyl, 4-methoxyphenyl, 3-methoxyphenyl, 3-chlorophenyl, 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 3',5'-dimethoxybiphenyl-4-yl, 2',4'-dimethoxybiphenyl-4-yl, biphenyl-4-yl, naphthalen-1-yl, naphthalen-2-yl or benzofuran-5-yl, phenylethyl, cyclohexyl, 2-furyl, thiophen-2-yl, thiophen-3-yl, 5-(3,4-dimethoxyphenyl)-thiophen-2-yl or 5-[2,2']bithiophenyl,
- n is 1 and
- m is 1.

The compounds of the formula I and also the starting substances for their preparation are otherwise prepared by methods known per se, such as are described in the literature (e.g. in the standard works such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), namely under reaction conditions which are known and suitable for the reactions mentioned. In this case, use can also be made of variants which are known per se, but not mentioned here in greater detail.

The starting substances, if desired, can also be formed in situ such that they are not isolated from the reaction mixture, but immediately reacted further to give the compounds of the formula I.

The compounds of the formula I can be obtained by liberating them from their functional derivatives by solvolysis, in particular hydrolysis or by hydrogenolysis.

Preferred starting substances for the solvolysis or hydrogenolysis are those which otherwise correspond to the formula I, but instead of one or more

free amino and/or hydroxyl groups contain corresponding protected amino and/or hydroxyl groups, in particular those which instead of an H-N- group carry an R'-N- group, in which R' is an amino protective group and/or those which instead of the H atom of a hydroxyl group carry a hydroxyl protective group, e.g. those which correspond to the formula I, but instead of a group -COOH carry a group -COOR", in which R" is a hydroxyl protective group.

A number of - identical or different - protected amino and/or hydroxyl groups can also be present in the molecule of the starting substance. If the protective groups present are different from one another, in many cases they can be removed selectively (lit.: T.W. Greene, P.G.M. Wuts, *Protective Groups in Organic Chemistry*, 2nd ed., Wiley, New York 1991, P.J. Kocienski, *Protecting Groups*, 1st ed. or Georg Thieme Verlag, Stuttgart - New-York, 1994).

The expression "amino protective group" is generally known and relates to groups which are suitable for protecting (for blocking) an amino group against chemical reactions, but which are easily removable after the desired chemical reaction has been carried out at other positions in the molecule. Typical groups of this type are, in particular, unsubstituted or substituted acyl, aryl, aralkoxymethyl or aralkyl groups. Since the amino protective groups are removed after the desired reaction (or reaction sequence), their nature and size is otherwise not critical; however, those having 1-20, in particular 1-8, C atoms are preferred. The expression "acyl group" is to be interpreted in the widest sense in connection with the present process. It includes acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids and, in particular, alkoxycarbonyl groups, aryloxycarbonyl groups and especially aralkoxycarbonyl groups. Examples of acyl groups of this type are alkanoyl such as acetyl, propionyl, butyryl; aralkanoyl such as phenylacetyl; aroyl

such as benzoyl or toluyl; aryloxyalkanoyl such as POA; alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, BOC, 2-iodoethoxycarbonyl; aralkyloxycarbonyl such as CBZ ("carbobenzoxy"), 4-methoxybenzyloxycarbonyl (MOZ), 4-Nitro-benzyloxycarbonyl oder 9-fluorenylmethoxycarbonyl (Fmoc); 2-(phenylsulfonyl)ethoxycarbonyl; trimethylsilylethoxycarbonyl (Teoc) or arylsulfonyl such as 4-methoxy-2,3,6-trimethylphenyl-sulfonyl (Mtr). Preferred amino protective groups are BOC, furthermore CBZ, Fmoc, benzy and acetyl; particularly preferred Fmoc.

The expression "hydroxyl protective group" is also generally known and relates to groups which are suitable for protecting a hydroxyl group against chemical reactions, but which are easily removable after the desired chemical reaction has been carried out at other positions in the molecule.

Typical groups of this type are the abovementioned unsubstituted or substituted aryl, aralkyl, aroyl or acyl groups, furthermore also alkylgroups, alkyl-, aryl- or aralkylsilylgroups or O,O- or O,S-acetals. The nature and size of the hydroxyl protective groups is not critical, since they are removed again after the desired chemical reaction or reaction sequence; groups having 1-20, in particular 1-10 C atoms, are preferred. Examples of hydroxyl protective groups are, inter alia, benzyl, 4-methoxybenzyl oder 2,4-dimethoxybenzyl, aroyl groups such as benzoyl or p-nitrobenzoyl, acyl groups such as acetyl or pivaloyl, p-toluolsulfonyl, alkyl groups such as methyl or tert-butyl, but also allyl, alkylsilyl groups such as trimethylsilyl (TMS), triisopropylsilyl (TIPS), tert-butyldimethylsilyl (TBS) or triethylsilyl, trimethylsilylethyl, aralkylsilyl groups such as tert-butyldiphenylsilyl (TBDPS), cyclic acetals such as isopropylidene-, cyclopentylidene-, cyclohexylidene-, benzyldiene-, p-methoxybenzyldiene- or o,p-dimethoxybenzyldieneacetal, acyclic acetals such as tetrahydropyranyl

(Thp), methoxymethyl (MOM), methoxyethoxymethyl (MEM), benzyloxymethyl (BOM) or methylthiomethyl (MTM). Acetyl, benzyl, tert-butyl or TBS being particularly preferred.

- 5 The liberation of the compounds of the formula I from their functional derivatives depending on the protective group used is known in the present literature such as T.W. Greene, P.G.M. Wuts, *Protective Groups in Organic Chemistry*, 2nd ed., Wiley, New York 1991, P.J. Kocienski, *Protecting Groups*, 1st ed., Georg Thieme Verlag, Stuttgart - New-York, 1994. In this
- 10 case, use can also be made of variants which are known per se, but not mentioned here in greater detail.

- The groups BOC and O-tert-butyl can preferably be removed, for example, using TFA in dichloromethane or using approximately 3 to 5N HCl in
- 15 dioxane at 15-30°C, the Fmoc group using an approximately 5 to 50% solution of dimethylamine, diethylamine or piperidine in DMF at 15-30°C.

- Preferred starting substances for the solvolysis or hydrogenolysis includes also those which otherwise correspond to the formula I, but are attached to
- 20 a solid phase. The liberation of the compounds of the formula I from the solid phase is known in the present literature such as Novabiochem - The Combinatorial Chemistry Catalog, March 99 and cited literature.

- The solid phase with a carbonate moiety as terminal functional group can
- 25 preferably be removed, for example, using TFA (50%) in dichloromethane.

- The quinazolinones of formula I can also preferably be prepared, using either solution or solid-phase techniques, by combining and reacting an anthranilic acid of formula II with an amine of formula III and if necessary
- 30 deprotect the given formula IV in which Q is then NH₂ or NHA and reacting

the compound of formul IV in which Q is NH_2 or NHA with an aldehyde of formula V.

As a rule, the starting compounds of the formulae II, III and V are known or commercially available.

5 The unknown compounds, however, can be prepared by methods known per se. The compounds of the formula II are anthranilic acids. It is furthermore possible to introduce appropriate substituents into the aromatic by conventional electrophilic or alternatively nucleophilic substitutions. Examples of Fmoc protected anthranilic acids, include, but are not limited to, Fmoc protected anthranilic acid, Fmoc protected 3-methyl anthranilic acid, Fmoc protected 3-methoxy anthranilic acid, Fmoc protected 3-chloro anthranilic acid or Fmoc protected 4-chloro anthranilic acid.

15 Solid-phase techniques may be employed to condense anthranilic acids of formula II and the amine component of formula III which is resin bound (R^2 or R^3 is solid phase).

The amines of formula III in which R^2 or R^3 are H, as a rule, are also commercially available and can be attached to the suitable resin by coupling procedures well known in the art and as described in the ensuing Examples. Furthermore, syntheses for the preparation of amines of formula 20 III, such as, for example, the Gabriel synthesis, can be used.

The aldehydes of formula V, as a rule, are also commercially available. Furthermore, syntheses for the preparation of aldehydes of formula V, such as, for example, the oxidation of an alcohol, can be used.

25 As a rule, the reactions and the attachment to the resin are carried out in an inert solvent. Depending on the conditions used, the reaction time is between a few minutes and a number of days, the reaction temperature between approximately 0° and 150°C , normally between 20° and 130°C .

30

Suitable inert solvents are, for example, hydrocarbons such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons such as trichloroethylene, 1,2-dichloroethane, carbon tetrachloride, chloroform or dichloromethane; alcohols such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers such as ethylene glycol monomethyl or monoethyl ether (methyl glycol or ethyl glycol), ethylene glycol dimethyl ether (diglyme); ketones such as acetone or butanone; amides such as acetamide, N-methylpyrrolidone (NMP), dimethylacetamide or dimethylformamide (DMF); nitriles such as acetonitrile; sulfoxides such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids such as formic acid or acetic acid; nitro compounds such as nitromethane or nitrobenzene; esters such as ethyl acetate or mixtures of the solvents mentioned.

The reaction of the compounds of formula II with compounds of formula III is analogously to the coupling of peptides. The condensation reaction of formula II with formula III is preferably carried out in an inert solvent as indicated above in the presence of a dehydrating agent, such as, dicyclohexylcarbodiimide (DCC), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide-hydrochlorid (EDC) or diisopropylcarbodiimide (DIC), further for instance in the presence of an anhydride of propanphosphonic acid (see Angew. Chem. 1980, 92, 129), diphenylphosphorylazide or 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline.

Particularly preferred is the presence of a coupling agent, such as TBTU (O-(benzotriazol-1-yl)-N,N,N',N'-bis-(tetramethylene)-uronium tetrafluoroborate) or O-(benzotriazol-1-yl)-N,N,N',N'-bis-(tetramethylene)-uronium hexafluorophosphate.

A compound of formula II in which X is a reactive esterified OH group can be synthesized by reacting a compound of formula II in which X is OH with HOBT (1-hydroxybenzotriazole) or N-hydroxysuccinimide (e.g. in the standard works such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart).

For the preparation of compounds of the formula I in which R^2 or R^3 are $-C(=NH)-NH-$, a compound of formula I in which R^2 and R^3 are H can be treated with an amidinating agent. The preferred amidinating agent is 1-amidino-3,5-dimethylpyrazole (DPFN), which is employed, in particular, in the form of its nitrate, or pyrazole-1-carboxamidine. The reaction is expediently carried out with addition of a base such as triethylamine or ethyldiisopropylamine in an inert solvent or solvent mixture, e.g. DMF at temperatures between 0° and 150°C , preferably between 60° and 120°C .

For the preparation of compounds of the formula I in which R^4 is unsubstituted or substituted biphenyl, 5-(3,4-dimethoxyphenyl)-thiophen-2-yl or 5-[2,2']bithiophenyl, an appropriate compound of the formula I in which R^4 is phenyl chloride, phenyl bromide, phenyl iodide, thiophenyl chloride, thiophenyl bromide or thiophenyl iodide can be reacted with the appropriate boronic acid derivatives in a Suzuki type coupling reaction. This reaction is expediently carried out under Palladium catalysis with different phosphines as coordination ligands, e.g. $\text{Pd}(\text{P}(\text{Ph})_3)_2$, $\text{Pd}(\text{II})\text{Cl}_2\text{dppf}$, $\text{PdOAc}_2 + \text{P}(\text{R}^*)_3$ ($\text{R}^* = \text{phenyl, cyclohexyl, tert-butyl}$) etc. in the presence of a base such as potassium carbonate, caesium carbonate, DBU, NaOH, in an inert solvent or solvent mixture, e.g. DMF or 1,4-dioxane at temperatures between 0° and 150° , preferably between 60° and 120° . Depending on the conditions used, the reaction time is between a few minutes and a number of days. The boronic acid derivatives can be prepared by conventional methods or are commercially available. The reactions can be carried out in analogy to

the methods indicated in Suzuki et al., J. Am. Chem. Soc. 1989, 111, 314ff., Suzuki et al., Chem. Rev. 1995, 95, 2457ff and G.C. Fu et al. Angew. Chem 1998, 110, 3586.

5 A base of the formula I can be converted into the associated acid addition salt using an acid, for example by reaction of equivalent amounts of the base and of the acid in an inert solvent such as ethanol and subsequent evaporation. Acids which give physiologically acceptable salts are particularly suitable for this reaction. Thus inorganic acids can be used, e.g.
10 sulfuric acid, nitric acid, hydrohalic acids such as hydrochloric acid or hydrobromic acid, phosphoric acids such as orthophosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic mono- or polybasic carboxylic, sulfonic or sulfuric acids, e.g. formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic
15 acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, p-toluenesulfonic acid, naphthalenemono- and disulfonic acids or laurylsulfuric acid. Salts with physiologically unacceptable acids, e.g.
20 picrates, can be used for the isolation and/or purification of the compounds of the formula I.

On the other hand, compounds of the formula I with bases (e.g. sodium or potassium hydroxide or carbonate) can be converted into the corresponding metal salts, in particular alkali metal or alkaline earth metal
25 salts, or into the corresponding ammonium salts.

The invention furthermore relates to pharmaceutical preparations comprising at least one compound of the formula I and/or one of its physiologically acceptable salts, which are prepared, in particular, in a
30 non-chemical way. In this case, the compounds of the formula I can be

brought into a suitable dose form together with at least one solid, liquid and/or semi-liquid excipient or auxiliary and, if appropriate, in combination with one or more other active compounds.

- 5 These preparations can be used as medicaments in human or veterinary medicine. Possible excipients are organic or inorganic substances which are suitable for enteral (e.g. oral) or parenteral administration or topical application and do not react with the novel compounds, for example water, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glyceryl triacetate, gelatin, carbohydrates such as lactose or starch, 10 magnesium stearate, talc and petroleum jelly. Tablets, pills, coated tablets, capsules, powders, granules, syrups, juices or drops are used, in particular, for oral administration, suppositories are used for rectal administration, solutions, preferably oily or aqueous solutions, furthermore suspensions, 15 emulsions or implants, are used for parenteral administration, and ointments, creams or powders are used for topical application. The novel compounds can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection preparations. The preparations indicated can be sterilized and/or can contain auxiliaries such as lubricants, 20 preservatives, stabilizers and/or wetting agents, emulsifiers, salts for affecting the osmotic pressure, buffer substances, colourants, flavourings and/or one or more other active compounds, e.g. one or more vitamins.

- 25 The compounds of the formula I and their physiologically acceptable salts act as adhesion receptor antagonists, in particular glycoprotein IbIX antagonists, and can be employed for the prophylaxis and/or therapy of thrombotic disorders and sequelae deriving therefrom. The disorders are acute coronary syndromes, angina pectoris, myocardial infarct, peripheral circulatory disorders, stroke, transient ischaemic attacks, arteriosclerosis 30 and reocclusion/restenosis after angioplasty/stent implantation.

In this case, the substances according to the invention are as a rule administered in the dose of the glycoprotein IIb/IIIa antagonist ReoPro® of preferably between approximately 1 and 500 mg, in particular between 5
5 and 100 mg, per dose unit. The daily dose is preferably between approximately 0.02 and 10 mg/kg of body weight. The specific dose for each patient depends, however, on all sorts of factors, for example on the efficacy of the specific compound employed, on the age, body weight, general state of health and sex, on the diet, on the time and route of
10 administration, and on the excretion rate, pharmaceutical combination and severity of the particular disorder to which the therapy applies. Oral administration is preferred.

Above and below, all temperatures are indicated in °C. In the following
15 examples, "customary working-up" for solution reactions means: if necessary, water is added, if necessary, depending on the constitution of the final product, the mixture is adjusted to pHs between 2 and 10 and extracted with ethyl acetate or dichloromethane, the organic phase is separated off, dried over sodium sulfate and evaporated, and the residue is
20 purified by chromatography on silica gel and/or by crystallization.

"Customary working-up" for solid-phase reactions means: the crude reaction is filtered and washed with DMF twice, then successively with methanol and methylene chloride three times, and finally once with methyl
25 tert-butyl ether. The resin is then dried in vacuo.


Mass spectrometry (MS) apparatuses Kratos Maldi III and Finnigan LCQ. (M+H)⁺ values or M⁺ values are determined.

EXAMPLES


Example 1:

3 grams (1.62 mmol) of p-nitrophenylcarbonate resin (1) [Novabiochem:
 5 0.54 mmol/g loading) is suspended in 30 ml of DMF then 8.1 mmol of C (3-Aminomethyl-cyclohexyl)-methylamine is added at room temperature. The reaction is then heated to 55° and left to stir for two days. The crude reaction is then customary worked up for solid-phase reactions affording the resin bound bis amine (2).

10




15



20

Analogously, by reaction of the p-nitrophenylcarbonate resin (1) with the bis amines of formula III



25

in which R² and R³ are H, excluding C-(3-aminomethyl-cyclohexyl)-methylamine, and n and m have the meanings indicated in Claim 1 the following resin bound bis amines are obtained:

cyclohexane-1,3-diamine, resin bound;

3-aminomethyl-cyclohexylamine, resin bound;

3-aminoethyl-cyclohexylamine, resin bound;

30

3-aminopropyl-cyclohexylamine, resin bound;

- C-(3-aminoethyl-cyclohexyl)-methylamine, resin bound;
C-(3-aminopropyl-cyclohexyl)-methylamine, resin bound;
C-(3-aminoethyl-cyclohexyl)-ethylamine, resin bound;
C-(3-aminopropyl-cyclohexyl)-propylamine, resin bound;
5 cyclohexane-1,4-diamine, resin bound;
4-aminomethyl-cyclohexylamine, resin bound;
4-aminoethyl-cyclohexylamine, resin bound;
4-aminopropyl-cyclohexylamine, resin bound;
C-(4-aminomethyl-cyclohexyl)-methylamine, resin bound;
10 C-(4-aminoethyl-cyclohexyl)-methylamine, resin bound;
C-(4-aminopropyl-cyclohexyl)-methylamine, resin bound;
C-(4-aminoethyl-cyclohexyl)-ethylamine, resin bound and
C-(4-aminopropyl-cyclohexyl)-propylamine, resin bound.


15 Example 2:

1. *Synthesis of Fmoc protected anthranilic acid*

29.15 mmol of anthranilic acid is taken in 100 ml of 1,4 dioxane then 145 mmol of sodium bicarbonate in 20 ml of water is added. Next, 32 mmol of Fmoc-Cl is added and the reaction is left to stir overnight at room
20 temperature. The reaction is then concentrated in vacuo and customary worked up for solution reactions. The resulting solid is triturated in ethyl ether affording the pure product.

2. *Coupling of Fmoc protected anthranilic acid to resin*

25 1 gram of resin (2) is suspended in 10 ml of DMF. The reaction is then treated with 1.62 mmol of Fmoc protected anthranilic acid, 1.62 mmol of HBTU, and 1.62 mmol of triethyl amine. The reaction is then allowed to shake overnight at room temperature. After customary working up, the resin is dried in vacuo affording resin bound anthranilic acid (3).




10 3. *Cleavage of Fmoc protected group*

1 gram resin (3) is suspended in 10 ml of 20% piperidine/DMF and shaken for 1.5 hours at room temperature. The reaction is then customary worked up for solid-phase reactions affording the free aniline (4).

15 4. *Aldehyde condensation and ring closure*

100 mg resin (4) is suspended in 1 ml of dimethyl acetamide then 200 μ l of acetic acid is added followed by the addition of 2.16 mmol of benzaldehyde. The reaction is then heated to 80° for two days. The reaction is then cooled to room temperature and customary worked up for solid-phase reactions affording the resin (5).

20




5. *Oxidation to quinazolinone*

100 mg resin (5) is suspended in 4 ml solution of 36 mg of DDQ in DMF. Then the reaction is allowed to shake overnight at room temperature. The

30

reaction is then customary worked up for solid-phase reactions affording quinazolinone (6) resin bound.




10 6. Cleavage of the final product 3-(3-aminomethyl-cyclohexylmethyl)-2-phenyl-3H-quinazolin-4-one

100 mg of resin (6) is suspended in 2 ml of a 50% trifluoroacetic acid/methylene chloride solution and shaken for 1.5 hours at room temperature. Customary working up for solid-phase reactions afforded 3-(3-aminomethyl-cyclohexylmethyl)-2-phenyl-3H-quinazolin-4-one;
 15 MS calc.: 347.4 found: 348.2.

Example 3:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa



cleavage of the Fmoc protecting group and reaction with benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-phenyl-3H-quinazolin-4-one;

MS calc.: 381.9 found: 382.2;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-phenyl-3H-quinazolin-4-one;

MS calc.: 361.5 found: 362.2;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-phenyl-3H-quinazolin-4-one;

MS calc.: 381.9 found: 382.2;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-phenyl-3H-quinazolin-4-one;

MS calc.: 377.5.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 2-methyl-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(2-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(2-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 375.5 found: 376.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(2-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(2-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 391.5 found: 392.2;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 361.5 found: 362.2.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-methyl-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(3-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(3-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 375.5 found: 376.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(3-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(3-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 391.5 found: 392.2;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(3-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 361.5 found: 362.2.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 4-methyl-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(4-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(4-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 375.5 found: 376.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(4-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(4-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 391.5 found: 392.2;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(4-methylphenyl)-3H-quinazolin-4-one;

MS calc.: 361.5 found: 362.2.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 4-tert-butyl-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(4-tert-butylphenyl)-3H-quinazolin-4-one;

MS calc.: 438.0 found: 438.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(4-tert-butylphenyl)-3H-quinazolin-4-one;

MS calc.: 417.6 found: 418.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(4-tert-butylphenyl)-3H-quinazolin-4-one;

MS calc.: 438.0 found: 438.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(4-tert-butylphenyl)-3H-quinazolin-4-one;

MS calc.: 433.6 found: 434.2;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(4-tert-butylphenyl)-3H-quinazolin-4-one;

MS calc.: 403.6 found: 404.3.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-chloro-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(3-chlorophenyl)-3H-quinazolin-4-one;

MS calc.: 416.4 found: 416.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(3-chlorophenyl)-3H-quinazolin-4-one;

MS calc.: 395.9 found: 396.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(3-chlorophenyl)-3H-quinazolin-4-one;

MS calc.: 416.3 found: 416.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(3-chlorophenyl)-3H-quinazolin-4-one;

MS calc.: 411.9 found: 412.1;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(3-chlorophenyl)-3H-quinazolin-4-one;

MS calc.: 381.9 found: 382.2.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 4-methoxy-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(4-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 411.9 found: 412.2;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(4-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 391.5 found: 392.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(4-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 411.9 found: 412.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(4-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 407.5 found: 408.2;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(4-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 377.5 found: 378.2.

20

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-methoxy-benzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(3-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 411.9 found: 412.1;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(3-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 391.5 found: 392.2;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(3-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 411.9 found: 412.2;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(3-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 407.5 found: 408.2;

15

with R' = H in formula IIa


3-(3-aminomethyl-cyclohexylmethyl)-2-(3-methoxyphenyl)-3H-quinazolin-4-one;

MS calc.: 377.5 found: 378.2.

20

Example 4:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa



cleavage of the Fmoc protecting group and reaction with 3,4,5-trimethoxybenzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

- 5 with R' = 3-Cl in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(3,4,5-trimethoxyphenyl)-3H-quinazolin-4-one;
- 10 with R' = 3-CH₃ in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(3,4,5-trimethoxyphenyl)-3H-quinazolin-4-one;
- 15 with R' = 4-Cl in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(3,4,5-trimethoxyphenyl)-3H-quinazolin-4-one;
- 20 with R' = 3-OCH₃ in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(3,4,5-trimethoxyphenyl)-3H-quinazolin-4-one;
- 25 with R' = H in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-2-(3,4,5-trimethoxyphenyl)-3H-quinazolin-4-one.
- 30 Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3,4-dimethoxybenzaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained
- 30 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(3,4-dimethoxyphenyl)-3H-quinazolin-4-one;

with R' = 3-CH₃ in formula IIa

5 3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(3,4-dimethoxyphenyl)-3H-quinazolin-4-one;

with R' = 4-Cl in formula IIa

10 3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(3,4-dimethoxyphenyl)-3H-quinazolin-4-one;

with R' = 3-OCH₃ in formula IIa

15 3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(3,4-dimethoxyphenyl)-3H-quinazolin-4-one;

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(3,4-dimethoxyphenyl)-3H-quinazolin-4-one.

20 Example 5:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with [2,2']bithiophenyl-5-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

25 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 470.1 found: 470.1;

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 449.6 found: 450.1;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 470.1 found: 470.1;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 465.6 found: 466.1;

15

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-3H-quinazolin-4-one;

MS calc.: 435.6 found: 436.1.

20

Example 6:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-furan-2-yl-propenal, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

30

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-(2-furan-2-yl-vinyl)-3H-quinazolin-4-one;

MS calc.: 397.9 found: 398.2;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-(2-furan-2-yl-vinyl)-3H-quinazolin-4-one;

MS calc.: 377.5 found: 378.3;

10 with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-(2-furan-2-yl-vinyl)-3H-quinazolin-4-one;

MS calc.: 397.9 found: 398.2;

15 with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-(2-furan-2-yl-vinyl)-3H-quinazolin-4-one;

MS calc.: 393.5 found: 394.3;

20 with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2-furan-2-yl-vinyl)-3H-quinazolin-4-one;

MS calc.: 363.5 found: 364.2.

25 Example 7:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with cyclohexanecarbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-cyclohexyl-3H-quinazolin-4-one;

MS calc.: 388.0 found: 388.2;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-cyclohexyl-3H-quinazolin-4-one;

MS calc.: 367.5 found: 368.3;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-cyclohexyl-3H-quinazolin-4-one;

MS calc.: 388.0 found: 388.2;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-cyclohexyl-3H-quinazolin-4-one;

MS calc.: 383.5 found: 384.3;

20

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-cyclohexyl-3H-quinazolin-4-one;

MS calc.: 353.5 found: 354.3.

25

Example 8:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-phenyl-propionaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-phenylethyl-3H-quinazolin-4-one;

MS calc.: 410.0 found: 410.3;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methyl-2-phenylethyl-3H-quinazolin-4-one;

MS calc.: 389.5 found: 390.4;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-phenylethyl-3H-quinazolin-4-one;

MS calc.: 410.0 found: 410.3;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-6-methoxy-2-phenylethyl-3H-quinazolin-4-one;

MS calc.: 405.5 found: 406.3;

20

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-phenylethyl-3H-quinazolin-4-one;

MS calc.: 375.5 found: 376.4.

25

Example 9:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with biphenyl-4-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-biphenyl-4-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 458.0 found: 458.2;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-biphenyl-4-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 437.6 found: 438.2;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-biphenyl-4-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 458.0 found: 458.2;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-biphenyl-4-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 453.6 found: 454.2;

20

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-biphenyl-4-yl-3H-quinazolin-4-one;

MS calc.: 423.6 found: 424.2.

25

Example 10:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with thiophene-3-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-3-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 387.9 found: 388.2;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-3-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 367.5 found: 368.2;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-3-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 387.9 found: 388.2;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-3-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 383.5 ; found: 384.2;

20

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-3-yl-3H-quinazolin-4-one;

MS calc.: 353.5 found: 354.2.

25

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with thiophene-2-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

- 3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-2-yl-6-chloro-3H-quinazolin-4-one;
MS calc.: 387.9 found: 388.2;
- 5 with R' = 3-CH₃ in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-2-yl-6-methyl-3H-quinazolin-4-one;
MS calc.: 367.5 found: 368.2;
- 10 with R' = 4-Cl in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-2-yl-7-chloro-3H-quinazolin-4-one;
MS calc.: 387.9 found: 388.1;
- 15 with R' = 3-OCH₃ in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-2-yl-6-methoxy-3H-quinazolin-4-one;
MS calc.: 383.5 found: 384.2;
- 20 with R' = H in formula IIa
3-(3-aminomethyl-cyclohexylmethyl)-2-thiophenyl-2-yl-3H-quinazolin-4-one;
MS calc.: 353.5 found: 354.2.
- Example 11:
- 25 Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with naphthalene-2-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained
- 30 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 432.0 found: 432.2;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 411.6 found: 412.2;

10 with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 432.0 found: 432.2;

15 with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 427.6 found: 428.2;

20 with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-3H-quinazolin-4-one;

MS calc.: 397.5 found: 398.2.

25 Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with naphthalene-1-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 432.0 found: 432.2;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 411.6 found: 412.2;

10 with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 432.0 found: 432.2;

15 with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 427.6 found: 428.2;

20 with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-3H-quinazolin-4-one;

MS calc.: 397.5 found: 398.2.

25 Example 12:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-phenyl-propenal, oxidation and cleavage from the solid phase, the following compounds are obtained

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-styryl-6-chloro-3H-quinazolin-4-one;

MS calc.: 407.9 found: 408.2;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-styryl-6-methyl-3H-quinazolin-4-one;

MS calc.: 387.5 found: 388.3;

with R' = 4-Cl in formula IIa

10 3-(3-aminomethyl-cyclohexylmethyl)-2-styryl-7-chloro-3H-quinazolin-4-one;

MS calc.: 407.9 found: 408.2;

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-styryl-6-methoxy-3H-quinazolin-4-

15 one;

MS calc.: 403.5 found: 404.3;

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-styryl-3H-quinazolin-4-one;

20 MS calc.: 373.5 found: 374.3.

Example 13:

25 Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with benzofuran-5-carbaldehyde, oxidation and cleavage from the solid phase, the following compounds are obtained

30 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-benzofuran-5-yl-6-chloro-3H-quinazolin-4-one;

MS calc.: 421.9 found: 422.2;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-benzofuran-5-yl-6-methyl-3H-quinazolin-4-one;

MS calc.: 401.5 found: 402.2;

10 with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-benzofuran-5-yl-7-chloro-3H-quinazolin-4-one;

MS calc.: 421.9 found: 422.2;

15 with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-benzofuran-5-yl-6-methoxy-3H-quinazolin-4-one;

MS calc.: 417.5 found: 418.1;

20 with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-benzofuran-5-yl-3H-quinazolin-4-one;

MS calc.: 387.5 found: 388.2.

25 Example 14:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-(4-dimethylamino-phenyl)-propenal, oxidation and cleavage from the solid phase, the following compounds are obtained

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(4-dimethylamino-phenyl)-vinyl]-6-chloro-3H-quinazolin-4-one;

MS calc.: 451.0;

5

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(4-dimethylamino-phenyl)-vinyl]-6-methyl-3H-quinazolin-4-one;

MS calc.: 430.6;

10

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(4-dimethylamino-phenyl)-vinyl]-7-chloro-3H-quinazolin-4-one;

MS calc.: 451.0;

15

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(4-dimethylamino-phenyl)-vinyl]-6-methoxy-3H-quinazolin-4-one;

MS calc.: 446.6;

20

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(4-dimethylamino-phenyl)-vinyl]-3H-quinazolin-4-one;

MS calc.: 416.6;

25

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 3-(2,5-dimethoxy-phenyl)-propenal, oxidation and cleavage from the solid phase, the following compounds are obtained

30

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(2,5-dimethoxy-phenyl)-vinyl]-6-chloro-3H-quinazolin-4-one;

5 with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(2,5-dimethoxy-phenyl)-vinyl]-6-methyl-3H-quinazolin-4-one;

with R' = 4-Cl in formula IIa

10 3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(2,5-dimethoxy-phenyl)-vinyl]-7-chloro-3H-quinazolin-4-one;

with R' = 3-OCH₃ in formula IIa

15 3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(2,5-dimethoxy-phenyl)-vinyl]-6-methoxy-3H-quinazolin-4-one;

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[2-(2,5-dimethoxy-phenyl)-vinyl]-3H-quinazolin-4-one.

20

Example 15:

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 4-bromo-benzaldehyde, Suzuki-reaction with 2,4-dimethoxyphenyl boronic acid as indicated afterwards, oxidation and cleavage from the solid phase, the following compounds are obtained

25

with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2',4'-dimethoxy-biphenyl-4-yl)-6-chloro-3H-quinazolin-4-one;

30

with R' = 3-CH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2',4'-dimethoxy-biphenyl-4-yl)-6-methyl-3H-quinazolin-4-one;

5

with R' = 4-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2',4'-dimethoxy-biphenyl-4-yl)-7-chloro-3H-quinazolin-4-one;

10

with R' = 3-OCH₃ in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(2',4'-dimethoxy-biphenyl-4-yl)-6-methoxy-3H-quinazolin-4-one;

with R' = H in formula IIa

15

3-(3-aminomethyl-cyclohexylmethyl)-2-(2',4'-dimethoxy-biphenyl-4-yl)-3H-quinazolin-4-one.

Suzuki reaction according to G.C. Fu et al., Angew. Chem. 1998, 110, 3586-3587:

20

1 gram of resin bound 3-(3-aminomethyl-cyclohexylmethyl)-2-(4-bromophenyl)-3H-quinazolin-4-one is suspended in 10 ml of 1,4-dioxane. The reaction is then treated with 1.62 mmol Cs₂CO₃, 1.62 mmol of 2,4-dimethoxyphenyl boronic acid and 10 mol% ([Pd₂(dba)₃] + P(tert-Bu)₃). The reaction is then allowed to shake at 80° until conversion is complete. After cooling the reaction mixture, it is worked up as is customary.

25

Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 4-bromo-benzaldehyde, Suzuki-reaction with 3,5-dimethoxyphenyl boronic

30

acid as indicated afterwards, oxidation and cleavage from the solid phase, the following compounds are obtained

with R' = 3-Cl in formula IIa

- 5 3-(3-aminomethyl-cyclohexylmethyl)-2-(3',5'-dimethoxy-biphenyl-4-yl)-6-chloro-3H-quinazolin-4-one;

with R' = 3-CH₃ in formula IIa

- 10 3-(3-aminomethyl-cyclohexylmethyl)-2-(3',5'-dimethoxy-biphenyl-4-yl)-6-methyl-3H-quinazolin-4-one;

with R' = 4-Cl in formula IIa

- 15 3-(3-aminomethyl-cyclohexylmethyl)-2-(3',5'-dimethoxy-biphenyl-4-yl)-7-chloro-3H-quinazolin-4-one;

with R' = 3-OCH₃ in formula IIa

- 3-(3-aminomethyl-cyclohexylmethyl)-2-(3',5'-dimethoxy-biphenyl-4-yl)-6-methoxy-3H-quinazolin-4-one;

- 20 with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-(3',5'-dimethoxy-biphenyl-4-yl)-3H-quinazolin-4-one.

- 25 Analogously to example 2, by reaction of resin (2) with a compound of formula IIa, cleavage of the Fmoc protecting group and reaction with 5-bromo-thiophenyl-2-carbaldehyde, Suzuki-reaction with 3,4-dimethoxyphenyl boronic acid as indicated afterwards, oxidation and cleavage from the solid phase, the following compounds are obtained

- 30 with R' = 3-Cl in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[5-(2,4-dimethoxy-phenyl)-2-thiophenyl]-6-chloro-3H-quinazolin-4-one;

with R' = 3-CH₃ in formula IIa

5 3-(3-aminomethyl-cyclohexylmethyl)-2-[5-(2,4-dimethoxy-phenyl)-2-thiophenyl]-6-methyl-3H-quinazolin-4-one;

with R' = 4-Cl in formula IIa

10 3-(3-aminomethyl-cyclohexylmethyl)-2-[5-(2,4-dimethoxy-phenyl)-2-thiophenyl]-7-chloro-3H-quinazolin-4-one;

with R' = 3-OCH₃ in formula IIa

15 3-(3-aminomethyl-cyclohexylmethyl)-2-[5-(2,4-dimethoxy-phenyl)-thiophen-2-yl]-6-methoxy-3H-quinazolin-4-one;

with R' = H in formula IIa

3-(3-aminomethyl-cyclohexylmethyl)-2-[5-(2,4-dimethoxy-phenyl)-thiophen-2-yl]-3H-quinazolin-4-one.

20 The following examples relate to pharmaceutical preparations:

Example A: Injection vials

A solution of 100 g of an active compound of the formula I and 5 g of disodium hydrogenphosphate is adjusted to pH 6.5 in 3 l of double-distilled water using 2N hydrochloric acid, sterile-filtered, dispensed into injection
25 vials, lyophilized under sterile conditions and aseptically sealed. Each injection vial contains 5 mg of active compound.

Example B: Suppositories

A mixture of 20 g of an active compound of the formula I is melted with 100 g of soya lecithin and 1400 g of cocoa butter, poured into moulds and allowed to cool. Each suppository contains 20 mg of active compound.

5

Example C: Solution

A solution is prepared from 1 g of an active compound of the formula I, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of double-distilled water. The mixture is adjusted to pH 6.8, made up to 1 l and sterilized by irradiation. This solution can be used in the form of eye drops.

10

Example D: Ointment

500 mg of an active compound of the formula I is mixed with 99.5 g of petroleum jelly under aseptic conditions.

15

Example E: Tablets

A mixture of 1 kg of active compound of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 g of talc and 0.1 kg of magnesium stearate is compressed in a customary manner to give tablets such that each tablet contains 10 mg of active compound.

20

Example F: Coated tablets

Analogously to Example E, tablets are pressed which are then coated with a coating of sucrose, potato starch, talc, tragacanth and colourant in a customary manner.

25

Example G: Capsules

2 kg of active compound of the formula I are dispensed into hard gelatin capsules in a customary manner such that each capsule contains 20 mg of the active compound.

30

Example H: Ampoules

A solution of 1 kg of active compound of the formula I in 60 ml of double-distilled water is sterile-filtered, dispensed into ampoules, lyophilized under sterile conditions and aseptically sealed. Each ampoule contains 10 mg of active compound.

10

15


20

25

30

What is claimed is:

1. Compounds of the formula I



in which

10 R and R¹ are independently of each other H, A, OH, OA, OCH₂-Ar, Hal, NH₂, NHA, NA₂, NO₂, CN, C(O)R², CONH₂, CONHA, CONA₂, COOH, COOA or SO₂A,

R² and R³ are independently of each other H, A, -C(=NH)-NH₂ or solid phase,

R⁴ is Ar, phenylalkyl, cycloalkyl or Het,

15 Y may be absent and, if present, is alkenyl having 2 to 4 carbon atoms,

A is unbranched or branched alkyl having 1 to 6 carbon atoms,

Ar is phenyl, naphthyl, biphenyl or benzofuranyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂,

20 Het is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, NH₂, NHA, NA₂, COOH, COOA, phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂, SO₂NH₂, SO₂NAH or SO₂NA₂ or thiophenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA,

25

30

CF₃, OCF₃, Hal, CN, COOH, COOA, NH₂, NHA, NA₂, NO₂,
SO₂NH₂, SO₂NAH or SO₂NA₂,

Hal is F, Cl, Br or I,

n is 0, 1, 2 or 3,

5 m is 0, 1, 2 or 3,

and their pharmaceutically tolerable salts and solvates.

2. Compounds of the formula I according to Claim 1


- 10 a) 3-(3-aminomethyl-cyclohexylmethyl)-2-[2,2']bithiophenyl-5-yl-6-methoxy-3H-quinazolin-4-one,
- b) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-6-methoxy-3H-quinazolin-4-one;
- c) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-6-methyl-3H-quinazolin-4-one;
- 15 d) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-1-yl-3H-quinazolin-4-one;
- e) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-6-methoxy-3H-quinazolin-4-one;
- f) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-3H-quinazolin-4-one;
- 20 g) 3-(3-aminomethyl-cyclohexylmethyl)-2-naphthalen-2-yl-6-methyl-3H-quinazolin-4-one;
- h) 3-(3-aminomethyl-cyclohexylmethyl)-6-chloro-2-naphthalen-2-yl-3H-quinazolin-4-one;
- 25 i) 3-(3-aminomethyl-cyclohexylmethyl)-7-chloro-2-naphthalen-2-yl-3H-quinazolin-4-one;
- and their physiologically acceptable salts and solvates.

3. Process for the preparation of the compounds of the formula I according to Claim 1 and their salts or solvates, characterized in that

30

a) a compound of the formula I is liberated from one of its functional derivatives by treating with a solvolysing or hydrogenolysing agent, or

b) in stage 1) a compound of the formula II




in which

10 X is Cl, Br, OH or a reactive esterified OH group and

Q is NH₂ or NHA, either of which is optionally protected, and


R and R¹ are optionally protected when they are or contain NH₂ or NHA,

is reacted with a compound of the formula III



in which R², R³, n and m have the meanings indicated in Claim 1,


to give a compound of formula IV



in which R, R¹, R², R³, Q, n and m have the meanings indicated above,

and

25 in stage 2) a compound of formula IV as indicated above is if necessary deprotected to give a compound of formula IV in which Q is NH₂ or NHA and is reacted with a compound of formula V



30 in which R⁴ and Y have the meanings indicated in Claim 1,

or

c) a radical R, R¹, R², R³ and/or R⁴ is converted into another radical R, R¹, R², R³ and/or R⁴ by, for example

- converting an amino group into a guanidino group by reaction with an amidinating agent,
- 5 - reducing a nitro group, sulfonyl group or sulfoxyl group,
- etherifying an OH group or subjecting an OA group to ether cleavage,
- alkylating a primary or secondary amino group,
- partially or completely hydrolysing a CN group,
- 10 - cleaving an ester group or esterifying a carboxylic acid radical,
- reacting an aryl bromide, aryl iodide, heteroaryl bromide or heteroaryliodide to give the corresponding coupling products by means of a Suzuki coupling with boronic acids,
- or carrying out a nucleophilic or electrophilic substitution,
- 15 and/or
- (e) a base or acid of the formula I is converted into one of its salts or solvates.

4. Compounds of the formula I according to Claim 1 and their
20 physiologically acceptable salts or solvates as pharmaceutical active compounds.

5. Compounds of the formula I according to Claim 1 and their
25 physiologically acceptable salts or solvates as glycoprotein IbIX antagonists.

6. Compounds of the formula I according to Claim 1 and their
30 physiologically acceptable salts or solvates as glycoprotein IbIX antagonists for the control of thrombotic disorders and sequelae deriving therefrom.

7. Pharmaceutical preparation characterized in that it contains at least one compound of the formula I according to Claim 4 and/or one of its physiologically acceptable salts or solvates.
- 5 8. Use of compounds of the formula I according to Claim 1 and/or their physiologically acceptable salts or solvates for the production of a pharmaceutical preparation for the control of thrombotic disorders and sequelae deriving therefrom or for use as anti-adhesive substances.
- 10 9. Use of compounds of the formula I according to Claim 4 and/or their physiologically acceptable salts or solvates for the production of a pharmaceutical preparation for the treatment of illnesses, such as for the prophylaxis and/or therapy of thrombotic disorders, as well as sequelae such as, for example, myocardial infarct, arteriosclerosis, angina
- 15 pectoris, acute coronary syndromes, peripheral circulatory disorders, stroke; transient ischaemic attacks, reocclusion/restenosis after angioplasty/stent implantations or as anti-adhesive substances for implants, catheters or heart pacemakers.
- 20
- 25
- 30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number
WO 01/23364 A1(51) International Patent Classification⁷: C07D 239/91,
405/04, 409/04, A61K 31/517, A61P 9/10

(21) International Application Number: PCT/EP00/08939

(22) International Filing Date:
13 September 2000 (13.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/407,939 28 September 1999 (28.09.1999) US(71) Applicant (for all designated States except US): MERCK
PATENT GMBH [DE/DE]; Frankfurter Strasse 250,
64293 Darmstadt (DE).

(72) Inventors; and


(75) Inventors/Applicants (for US only): MEDERSKI,
Werner [DE/DE]; Am Ohlenberg 29, 64390 Erzhäusen
(DE). DEVANT, Ralf [DE/DE]; Frankfurter Land-
strasse 135, 64291 Darmstadt (DE). BARNICKEL,
Gerhard [DE/DE]; Emilstrasse 27, 64298 Darmstadt
(DE). BERNOTAT-DANIELOWSKI, Sabine [DE/DE];
Liebigstrasse 5, 61231 Bad Nauheim (DE). MELZER,
Guido [DE/DE]; Mörikestrasse 6, 65719 Hofheim/Ts.
(DE). DHANOA, Daljit [US/US]; 2037 Skiles Blvd., West
Chester, PA 19382 (US). ZHAO, Bao-Ping [CN/US];3606 Hunters Glen Drive, Plainsboro, NJ 08536 (US).
RINKER, James [US/US]; 1115 Fern Avenue, Kenhurst,
PA 19607 (US). PLAYER, Mark [US/US]; 5022 Swan
Drive, Phoenixville, PA 1946 (US). SOLL, Richard
[US/US]; 324 Glenn Avenue, Lawrenceville, NJ 08648
(US).(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: QUINAZOLINONES



(57) Abstract: Quinazolinones of formula (I) in which R, R¹, R², R³, R⁴, Y, n and m have the meaning indicated in Patent Claim 1, and their salts or solvates as glycoprotein IbIX antagonists.

Docket No.
Merck 2032 A

Declaration and Power of Attorney For Patent Application

English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Quinazolinones

the specification of which

(check one)

☐
☒

is attached hereto.

was filed on 13.09.2000 as United States Application No. or PCT International
Application Number PCT/EP00/08939
and was amended on _____

(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

99-407,939
(Number)

USA
(Country)

28.09.1999
(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

_____	_____
(Application Serial No.)	(Filing Date)

_____	_____
(Application Serial No.)	(Filing Date)

_____	_____
(Application Serial No.)	(Filing Date)

I hereby claim the benefit under 35 U.S.C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States of PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112. I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C.F.R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

_____	_____	_____
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

_____	_____	_____
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

_____	_____	_____
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint

I. William Millen (Reg. No. 19,544)
 John L. White (Reg. No. 17,746)
 Anthony J. Zelano (Reg. No. 27,969)
 Alan E. J. Branigan (Reg. No. 20,565)
 John R. Moses (Reg. No. 24,983)
 Harry B. Shubin (Reg. No. 32,004)
 Brion P. Heaney (Reg. No. 32,542)
 Richard J. Traverso (Reg. No. 30,595)

John A. Sopp (Reg. No. 33,103)
 Richard M. Lebovitz (Reg. No. 37,067)
 John H. Thomas (Reg. No. 33,460)
 Catherine M. Joyce (Reg. No. 40,668)
 James T. Moore (Reg. No. 35,619)
 James E. Ruland (Reg. No. 37,432)
 Nancy Axelrod (Reg. No. 44,014)
 Jennifer J. Branigan (Reg. No. 40,921)

to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Send Correspondence to: MILLEN; WHITE; ZELANO & BRANIGAN, P.C.
 Arlington Courthouse Plaza I
 2200 Clarendon Blvd., Suite 1400
 Arlington, VA 22201

Direct Telephone Calls to: (name and telephone number)

1-00

Full name of sole or first inventor	27 MAI. 2002
<u>MEDERSKI, Werner</u>	Date
Sole or first inventor's	
<i>Werner Mederski</i>	
Residence	
<u>Katzeneinbogenweg 164673 Zwingenberg, Germany</u>	DEX
Citizenship	
<u>German</u>	
Post Office Address	
<u>c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany</u>	

2-00

Full name of second inventor, if any	27 MAI. 2002
<u>DEVANT, Ralf</u>	Date
Second inventor's signature	
<i>Devant, Ralf</i>	
Residence	
<u>Frankfurter Landstrasse 135, 64291 Darmstadt, Germany</u>	DEX
Citizenship	
<u>German</u>	
Post Office Address	
<u>c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany</u>	

3-00

Full name of third inventor, if any	27 MAI. 2002
BARNICKEL, Gerhard	Date
Third inventor's signature <i>Gerhard Barnickel</i>	
Residence	
Emilstrasse 27, 64293 Darmstadt, Germany	DEX
Citizenship	
German	
Post Office Address	
c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany	

4-00

Full name of fourth inventor, if any	27 MAI. 2002
BERNOTAT-DANIELOWSKI, Sabine	Date
Fourth inventor's signature <i>Sabine Bernotat-Danowski</i>	
Residence	
Liebigstrasse 5, 61231 Bad Nauheim, Germany	DEX
Citizenship	
German	
Post Office Address	
c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany	

5-00

Full name of fifth inventor, if any	27 MAI. 2002
MELZER, Guido	Date
Fifth inventor's signature <i>Guido Melzer</i>	
Residence	
Moerikestrasse 6, 65719 Hofheim/Taunus, Germany	DEX
Citizenship	
Austrian	
Post Office Address	
c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany	

6-00

Full name of sixth inventor, if any	27 MAI. 2002
CEZANNE, Bertram	Date
Sixth inventor's signature <i>Bertram Cezanne</i>	
Residence	
Bahnstrasse 74, 64546 Mörfelden-Walldorf, Germany	DEX
Citizenship	
German	
Post Office Address	
c/o Merck KGaA, Frankfurter Str. 250, 64293 Darmstadt, Germany	

7-DD

Full name of seventh inventor, if any	
DHANO, Daljit	27 MAY 2002
seventh inventor's signature	Date
<i>Daljit Dhano</i>	
Residence	
13794 Boquita Dr., Del Mar, CA 92014, USA CD	
Citizenship	
USA	
Post Office Address	

Full name of eighth inventor, if any	
ZHAO, Bao-Ping	
Eighth inventor's signature	Date
Residence	
5 Windsor Pond Rd., West Windsor, NJ 08550, USA	
Citizenship	
CA	
Post Office Address	

Full name of ninth inventor, if any	
RINKER, James	
Ninth inventor's signature	Date
Residence	
1115 Fern Ave., Kenhorst, PA 19607, USA	
Citizenship	
USA	
Post Office Address	

Full name of tenth inventor, if any	
PLAYER, Mark	
Tenth inventor's signature	Date
Residence	
5022 Swan Dr., Phoenixville, Pa 19607, USA	
Citizenship	
USA	
Post Office Address	

Full name of seventh inventor, if any	
DHANOA; Daljit	
seventh inventor's signature	Date
Residence	
13794 Boquita Dr., Del Mar, CA 92014; USA	
Citizenship	
USA	
Post Office Address	

8-00

Full name of eighth inventor, if any		27 MAY. 2002
ZHAO, Bao-Ping		Date
Eighth inventor's signature		
Residence		
5 Windsor Pond Rd., West Windsor, NJ 08550, USA NJ.		
Citizenship		
CA		
Post Office Address		

9-00

Full name of ninth inventor, if any		27 MAY. 2002
RINKER, James		Date
Ninth inventor's signature		
Residence		
1115 Fern Ave., Kenhorst, PA 19607, USA PA.		
Citizenship		
USA		
Post Office Address		

10-00

Full name of tenth inventor, if any		27 MAY. 2002
PLAYER, Mark		Date
Tenth inventor's signature		
Residence		
5022 Swan Dr., Rhoenixville, Pa 19607, USA PA.		
Citizenship		
USA		
Post Office Address		

11-62

Full name of eleventh inventor, if any	
SOLL; Richard	27 MAY. 2002
Eleventh inventor's signature	Date
Residence	
324 Glenn Ave., Lawrencehill, NJ 08648, USA	
Citizenship	
USA	
Post Office Address	

Full name of fourth inventor, if any	
Fourth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

Full name of fifth inventor, if any	
Fifth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	